



Swine Flu

What is Swine Influenza?

Swine Influenza (swine flu) is a respiratory disease of pigs caused by type A influenza virus that regularly causes outbreaks of influenza in pigs. Swine flu viruses cause high levels of illness and low death rates in pigs. Swine influenza viruses may circulate among swine throughout the year, but most outbreaks occur during the late fall and winter months similar to outbreaks in humans. The classical swine flu virus (an influenza type A H1N1 virus) was first isolated from a pig in 1930.

How many swine flu viruses are there?

Like all influenza viruses, swine flu viruses change constantly. Pigs can be infected by avian influenza and human influenza viruses as well as swine influenza viruses. When influenza viruses from different species infect pigs, the viruses can reassort (i.e. swap genes) and new viruses that are a mix of swine, human and/or avian influenza viruses can emerge. Over the years, different variations of swine flu viruses have emerged. At this time, there are four main influenza type A virus subtypes that have been isolated in pigs: H1N1, H1N2, H3N2, and H3N1. However, most of the recently isolated influenza viruses from pigs have been H1N1 viruses.

Can humans catch swine flu?

Swine flu viruses do not normally infect humans. However, sporadic human infections with swine flu have occurred. Most commonly, these cases occur in persons with direct exposure to pigs (e.g. children near pigs at a fair or workers in the swine industry). In addition, there have been documented cases of one person spreading swine flu to others. For example, an outbreak of apparent swine flu infection in pigs in Wisconsin in 1988 resulted in multiple human infections, and, although no community outbreak resulted, there was antibody evidence of virus transmission from the patient to healthcare workers who had close contact with the patient.

What happens?

The symptoms of swine flu in people are **similar to the symptoms of seasonal flu** in humans and may include:

- Fever (greater than 100°F or 37.8°C)
- Sore throat
- Cough
- Stuffy nose
- Chills
- Headache and body aches
- Fatigue
- Some people have reported diarrhea and vomiting associated with swine flu.
- Severe illness (pneumonia and respiratory failure) and deaths have been reported with swine flu infection in people.

TRAVEL VACCINATION CENTER

Somerset Professional Plaza, 1527, Rt. 27 South, Suite 1600, Somerset NJ-08873-3979

Dayton Professional Plaza, 401 Ridge road, Suite # 2, Dayton, NJ 08810

Tel: (732) 418- 1700 Fax: (732) 249 -9599



- Like seasonal flu, swine flu may cause a worsening of underlying chronic medical conditions.

Can people catch swine flu from eating pork?

No. Swine influenza viruses are not transmitted by food. You can not get swine influenza from eating pork or pork products. Eating properly handled and cooked pork and pork products are safe. Cooking pork to an internal temperature of 160°F kills the swine flu virus as it does other bacteria and viruses.

How does swine flu spread?

Spread of this swine influenza A (H1N1) virus is thought to be happening in the same way that seasonal flu spreads. Flu viruses are spread mainly from person to person through coughing or sneezing of people with influenza. Sometimes people may become infected by touching something with flu viruses on it and then touching their mouth or nose.

How can someone with the flu infect someone else?

Infected people may be able to infect others beginning 1 day before symptoms develop and up to 7 or more days after becoming sick. That means that you may be able to pass on the flu to someone else before you know you are sick, as well as while you are sick.

How can human infections with swine influenza be diagnosed?

To diagnose swine influenza A infection, a respiratory specimen would generally need to be collected within the first 4 to 5 days of illness (when an infected person is most likely to be shedding virus). However, some persons, especially children, may shed virus for 10 days or longer. Identification as a swine flu influenza A virus requires sending the specimen to CDC for laboratory testing.

Is the H1N1 swine flu virus the same as human H1N1 viruses?

No. The H1N1 swine flu viruses are antigenically very different from human H1N1 viruses and, therefore, vaccines for human seasonal flu would not provide protection from H1N1 swine flu viruses.

How long can an infected person spread swine flu to others?

People with swine influenza virus infection should be considered potentially contagious as long as they are symptomatic and possible for up to 7 days following illness onset. Children, especially younger children, might potentially be contagious for longer periods.

What surfaces are most likely to be sources of contamination?

Germs can be spread when a person touches something that is contaminated with germs and then touches his or her eyes, nose, or mouth. Droplets from a cough or sneeze of an infected person move through the air. Germs can be spread when a person touches respiratory droplets from another person on a surface like a desk and then touches their own eyes, mouth or nose before washing their hands.

How long can viruses live outside the body?

We know that some viruses and bacteria can live 2 hours or longer on surfaces like cafeteria tables, doorknobs, and desks. Frequent hand washing will help you reduce the chance of getting contamination from these common surfaces.

TRAVEL VACCINATION CENTER

Somerset Professional Plaza, 1527, Rt. 27 South, Suite 1600, Somerset NJ-08873-3979

Dayton Professional Plaza, 401 Ridge road, Suite # 2, Dayton, NJ 08810

Tel: (732) 418- 1700 Fax: (732) 249 -9599



How can we prevent the spread?

- **Avoid contact** with ill persons.
- When you cough or sneeze, **cover your nose and mouth** with a tissue or your sleeve (if you do not have a tissue). Throw used tissues in a trash can.
- After you cough or sneeze, wash your hands **with soap and water**, or use an **alcohol-based hand gel**.
- **If you think you are ill with flu**, avoid close contact with others as much as possible. **Stay at home or in your hotel room. Seek medical care if you are severely ill (such as having trouble breathing). There are antiviral medications for prevention and treatment of swine flu that a doctor can prescribe. Do not go to work, school, or travel while ill.**

What should I do to keep from getting the flu?

First and most important: wash your hands. Try to stay in good general health. Get plenty of sleep, be physically active, manage your stress, drink plenty of fluids, and eat nutritious food. Try not touch surfaces that may be contaminated with the flu virus. Avoid close contact with people who are sick.

Are there medicines to treat swine flu?

Yes. CDC recommends the use of oseltamivir or zanamivir for the treatment and/or prevention of infection with these swine influenza viruses. Antiviral drugs are prescription medicines (pills, liquid or an inhaler) that fight against the flu by keeping flu viruses from reproducing in your body. If you get sick, antiviral drugs can make your illness milder and make you feel better faster. They may also prevent serious flu complications. For treatment, antiviral drugs work best if started soon after getting sick (within 2 days of symptoms).

What should I do if I get sick?

Contact their health care provider, particularly if you are worried about your symptoms. Your health care provider will determine whether influenza testing or treatment is needed.

If you are sick, you should stay home and avoid contact with other people as much as possible to keep from spreading your illness to others.

If you become ill and experience any of the following warning signs, seek emergency medical care.

In children emergency warning signs that need urgent medical attention include:

- Fast breathing, trouble breathing, Difficulty breathing or shortness of breath
- Bluish skin color
- Not drinking enough fluids
- Not waking up or not interacting
- Being so irritable that the child does not want to be held
- Flu-like symptoms improve but then return with fever and worse cough
- Fever with a rash
- Pain or pressure in the chest or abdomen

TRAVEL VACCINATION CENTER

Somerset Professional Plaza, 1527, Rt. 27 South, Suite 1600, Somerset NJ-08873-3979

Dayton Professional Plaza, 401 Ridge road, Suite # 2, Dayton, NJ 08810

Tel: (732) 418- 1700 Fax: (732) 249 -9599



- Sudden dizziness
- Confusion
- Severe or persistent vomiting

TRAVEL VACCINATION CENTER

Somerset Professional Plaza, 1527, Rt. 27 South, Suite 1600, Somerset NJ-08873-3979

Dayton Professional Plaza, 401 Ridge road, Suite # 2, Dayton, NJ 08810

Tel: (732) 418- 1700 Fax: (732) 249 -9599