



## **Tetanus ad Diphtheria Vaccine (Td)**

Tetanus (lockjaw) and diphtheria are serious diseases. Tetanus is caused by a germ that enters the body through a cut or wound. Diphtheria spreads when germs pass from an infected person to the nose or throat of others.

Tetanus causes serious, painful spasms of all muscles.

It can lead to:

- "Locking" of the jaw so the patient cannot open his or her mouth or swallow

Diphtheria causes a thick coating in the nose, throat, or airway. It can lead to:

- Breathing problems
- Heart failure
- Paralysis
- Death

Vaccination is the best way to protect against tetanus and diphtheria, Because of vaccination; there are many fewer cases of these diseases. Cases are rare in children because most get DTP (Diphtheria, Tetanus, and Pertussis),

### **When should you get Td vaccine?**

Td is made for people 7 years of age and older.

People who have not gotten at least 3 doses of any tetanus and diphtheria vaccine (DTP, DTaP or DT) during their lifetime should do so using Td. After a person gets the third dose, a Td dose is needed every 10 years all through life.

Tell us if the person getting the vaccine:

- Ever had a serious allergic reaction or other problem with Td, or any other tetanus and diphtheria vaccine (DTP, DTaP or DT)
- Now has a moderate or severe illness
- Is pregnant

### **What are the risks from Td vaccine?**

---

#### **TRAVEL VACCINATION CENTER**

**1553 Ruth Road Suite 1, North Brunswick, NJ-08902**

**Dayton Professional Center, 401 Ridge Rd., Suite 2, Dayton, NJ 08810**

**Highland Park Pediatrics, 85 Raritan Ave, #410, Highland Park, NJ-08904**

**Tel: (732) 418- 1700 Fax: (732) 940-9700**



# **Central Jersey Pediatrics**

*Highland park Pediatrics*  
*Infants, Children & Adolescents*  
*www.cjpediatrics.com*

---

As with any medicine, there are very small risks that serious problems, even death, could occur after getting a vaccine.

The risks from the vaccine are much smaller than the risks from the diseases if people stopped using vaccine.

Almost all people who get Td have no problems from it.

## **Mild problems**

If these problems occur, they usually start within hours to a day or two after vaccination. They may last 1-2 days:

- Soreness, redness, or swelling where the shot was given. Acetaminophen or ibuprofen (non-aspirin) may be used to reduce soreness.

## **Severe problems**

These problems happen very rarely:

- Serious allergic reaction
- Deep, aching pain and muscle wasting in upper arm(s). This starts 2 days to 4 weeks after the shot, and may last many months.

## **What to do if there is a serious reaction:**

- \* Call a doctor or get the person to a doctor right away,
- \* Ask us or health department to file a Vaccine Adverse Event Report form or call: (800) 822-7967 (toll-free)

The National Vaccine Injury Compensation Program gives compensation (payment) for persons thought to be injured by vaccines. For details call: (800) 338-2382 (toll-free)

---

### **TRAVEL VACCINATION CENTER**

**1553 Ruth Road Suite 1, North Brunswick, NJ-08902**  
**Dayton Professional Center, 401 Ridge Rd., Suite 2, Dayton, NJ 08810**  
**Highland Park Pediatrics, 85 Raritan Ave, #410, Highland Park, NJ-08904**  
**Tel: (732) 418- 1700 Fax: (732) 940-9700**